

Bird Observations

Period: 2004-03-22 to 2004-04-18

Page 1 of 7

Species	Obs. days	Obs. %
Long-tailed Cormorant (Phalacrocorax africanus)	7	26%
Cattle Egret (Bubulcus ibis)	23	85%
Little Egret (Egretta garzetta)	3	11%
Yellow-billed Egret (Egretta intermedia)	4	15%
Great White Egret (Egretta alba)	1	4%
Grey Heron (Ardea cinerea)	1	4%
Black-headed Heron (Ardea melanocephala)	6	22%
Hamerkop (Scopus umbretta)	4	15%
Abdim's Stork (Ciconia abdimii)	2	7%
White-faced Duck (Dendrocygna viduata)	1	4%
Honey Buzzard (Pernis apivorus)	1	4%
Black-shouldered Kite (Elanus caeruleus)	11	41%
Yellow-billed Kite (Milvus migrans)	24	89%
Hooded Vulture (Necrosyrtes monachus)	25	93%
Gymnogene (Polyboroides typus)	1	4%
Eurasian Marsh Harrier (Circus aeruginosus)	1	4%
Montagu's Harrier (Circus pygargus)	3	11%
RINGTAIL ()	1	4%
Gabar Goshawk (Melierax gabar)	2	7%
African Goshawk (Accipiter tachiro)	1	4%
Shikra (Accipiter badius)	6	22%
Lizard Buzzard (Kaupifalco monogrammicus)	4	15%
Red-necked Buzzard (Buteo auguralis)	14	52%
Long-crested Eagle (Lophaetus occipitalis)	1	4%
Common Kestrel (Falco tinnunculus)	12	44%
Fox Kestrel (Falco alopex)	15	56%
Grey Kestrel (Falco ardosiaceus)	3	11%
Red-necked Falcon (Falco chicquera)	1	4%
European Hobby (Falco subbuteo)	2	7%
Lanner Falcon (Falco biarmicus)	25	93%
Double-spurred Francolin (Francolinus bicalcaratus)	18	67%
Common Quail (Coturnix coturnix)	2	7%
Stone Partridge (Ptilopachus petrosus)	25	93%

Database made by Turgor Henrik Dahl AB, SWEDEN

Species	Obs. days	Obs. %
Little Buttonquail (<i>Turnix sylvatica</i>)	1	4%
Black Crake (<i>Amaurornis flavirostris</i>)	1	4%
Painted Snipe (<i>Rostratula benghalensis</i>)	1	4%
Senegal Thick-knee (<i>Burhinus senegalensis</i>)	1	4%
Spotted Dikkop (<i>Burhinus capensis</i>)	1	4%
Temminck's Courser (<i>Cursorius temminckii</i>)	3	11%
Kittlitz's Plover (<i>Charadrius pecuarius</i>)	1	4%
Forbes's Plover (<i>Charadrius forbesi</i>)	1	4%
Senegal Wattled Plover (<i>Vanellus senegallus</i>)	7	26%
Spur-winged Plover (<i>Vanellus spinosus</i>)	8	30%
Common Snipe (<i>Gallinago gallinago</i>)	1	4%
Greenshank (<i>Tringa nebularia</i>)	4	15%
Wood Sandpiper (<i>Tringa glareola</i>)	3	11%
Common Sandpiper (<i>Actitis hypoleucos</i>)	4	15%
Speckled Pigeon (<i>Columba guinea</i>)	6	22%
Adamawa Turtle Dove (<i>Streptopelia hypopyrrha</i>)	25	93%
Laughing Dove (<i>Streptopelia senegalensis</i>)	25	93%
Vinaceous Dove (<i>Streptopelia vinacea</i>)	22	81%
Red-eyed Dove (<i>Streptopelia semitorquata</i>)	1	4%
Black-billed Dove (<i>Turtur abyssinicus</i>)	25	93%
Namaqua Dove (<i>Oena capensis</i>)	2	7%
Bruce's Green Pigeon (<i>Treron waalia</i>)	12	44%
Senegal Parrot (<i>Poicephalus senegalus</i>)	20	74%
Red-headed Lovebird (<i>Agapornis pullarius</i>)	5	19%
Violet Turaco (<i>Musophaga violacea</i>)	23	85%
Western Grey Plantain-eater (<i>Crinifer piscator</i>)	24	89%
Great Spotted Cuckoo (<i>Clamator glandarius</i>)	1	4%
Jacobin Cuckoo (<i>Clamator jacobinus</i>)	1	4%
Red-chested Cuckoo (<i>Cuculus solitarius</i>)	2	7%
African Grey Cuckoo (<i>Cuculus gularis</i>)	8	30%
Klaas's Cuckoo (<i>Chrysococcyx klaas</i>)	3	11%
Didric Cuckoo (<i>Chrysococcyx caprius</i>)	2	7%
Senegal Coucal (<i>Centropus senegalensis</i>)	24	89%
Spotted Eagle Owl (<i>Bubo africanus</i>)	20	74%
Marsh Owl (<i>Asio capensis</i>)	2	7%

Database made by Turgor Henrik Dahl AB, SWEDEN

Species	Obs. days	Obs. %
Freckled Rock Nightjar (<i>Caprimulgus tristigma</i>)	18	67%
Long-tailed Nightjar (<i>Caprimulgus climacurus</i>)	19	70%
Standard-winged Nightjar (<i>Macrodipteryx longipennis</i>	1	4%
African Palm Swift (<i>Cypsiurus parvus</i>)	25	93%
Alpine Swift (<i>Apus melba</i>)	1	4%
Mottled Swift (<i>Apus aequatorialis</i>)	1	4%
European Swift (<i>Apus apus</i>)	2	7%
Little Swift (<i>Apus affinis</i>)	20	74%
African White-rumped Swift (<i>Apus caffer</i>)	11	41%
Speckled Mousebird (<i>Colius striatus</i>)	25	93%
Malachite Kingfisher (<i>Alcedo cristata</i>)	4	15%
Pygmy Kingfisher (<i>Ceyx pictus</i>)	15	56%
Grey-headed Kingfisher (<i>Halcyon leucocephala</i>)	9	33%
Blue-breasted Kingfisher (<i>Halcyon malimbica</i>)	1	4%
Striped Kingfisher (<i>Halcyon chelicuti</i>)	25	93%
Pied Kingfisher (<i>Ceryle rudis</i>)	2	7%
Little Bee-eater (<i>Merops pusillus</i>)	11	41%
Red-throated Bee-eater (<i>Merops bulocki</i>)	25	93%
European Bee-eater (<i>Merops apiaster</i>)	21	78%
Northern Carmine Bee-eater (<i>Merops nubicus</i>)	1	4%
Abyssinian Roller (<i>Coracias abyssinica</i>)	1	4%
Red-billed Wood Hoopoe (<i>Phoeniculus purpureus</i>)	18	67%
Black Wood Hoopoe (<i>Rhinopomastus aterrimus</i>)	3	11%
Hoopoe (<i>Upupa epops</i>)	1	4%
Red-billed Hornbill (<i>Tockus erythrorhynchus</i>)	10	37%
African Grey Hornbill (<i>Tockus nasutus</i>)	25	93%
Yellow-fronted Tinkerbird (<i>Pogoniulus chrysoconus</i>)	25	93%
Vieillot's Barbet (<i>Lybius vielloti</i>)	15	56%
White-headed Barbet (<i>Lybius leucocephalus</i>)	4	15%
Bearded Barbet (<i>Lybius dubius</i>)	13	48%
Greater Honeyguide (<i>Indicator indicator</i>)	22	81%
Lesser Honeyguide (<i>Indicator minor</i>)	6	22%
European Wryneck (<i>Jynx torquilla</i>)	1	4%
Fine-spotted Woodpecker (<i>Campethera punctuligera</i>)	1	4%
Cardinal Woodpecker (<i>Dendropicos fuscescens</i>)	4	15%

Database made by Turgor Henrik Dahl AB, SWEDEN

Species	Obs. days	Obs. %
Grey Woodpecker (<i>Mesopicos goertae</i>)	18	67%
Rufous-naped Lark (<i>Mirafrā africana</i>)	2	7%
Sun Lark (<i>Galerida modesta</i>)	21	78%
Crested Lark (<i>Galerida cristata</i>)	9	33%
Chestnut-backed Sparrow-Lark (<i>Eremopterix leucotis</i>)	9	33%
Fanti Saw-wing (<i>Psalidoprocne obscura</i>)	10	37%
Lesser Striped Swallow (<i>Hirundo abyssinica</i>)	24	89%
Red-rumped Swallow (<i>Hirundo daurica</i>)	23	85%
African Rock Martin (<i>Hirundo fuligula</i>)	25	93%
Ethiopian Swallow (<i>Hirundo aethiopica</i>)	22	81%
European Swallow (<i>Hirundo rustica</i>)	17	63%
House Martin (<i>Delichon urbica</i>)	11	41%
Yellow Wagtail (<i>Motacilla flava</i>)	5	19%
Long-billed Pipit (<i>Anthus similis</i>)	4	15%
Plain-backed Pipit (<i>Anthus leucophrys</i>)	4	15%
Tree Pipit (<i>Anthus trivialis</i>)	11	41%
Red-throated Pipit (<i>Anthus cervinus</i>)	3	11%
Yellow-throated Longclaw (<i>Macronyx croceus</i>)	22	81%
Red-shouldered Cuckoo-shrike (<i>Campephaga</i>	10	37%
Yellow-throated Leaflove (<i>Chlorocichla flavicollis</i>)	25	93%
Common Bulbul (<i>Pycnonotus barbatus</i>)	25	93%
West African Thrush (<i>Turdus pelios</i>)	25	93%
Snowy-crowned Robin-Chat (<i>Cossypha niveicapilla</i>)	9	33%
White-crowned Robin-Chat (<i>Cossypha albicapilla</i>)	3	11%
Whinchat (<i>Saxicola rubetra</i>)	15	56%
Familiar Chat (<i>Cercomela familiaris</i>)	25	93%
Northern Ant-eating Chat (<i>Myrmecocichla aethiops</i>)	6	22%
Mocking Chat (<i>Myrmecocichla cinnamomeiventris</i>)	25	93%
African Moustached Warbler (<i>Melocichla mentalis</i>)	10	37%
Reed Warbler (<i>Acrocephalus scirpaceus</i>)	4	15%
Olivaceous Warbler (<i>Hippolais pallida</i>)	1	4%
Icterine Warbler (<i>Hippolais icterina</i>)	4	15%
Melodious Warbler (<i>Hippolais polyglotta</i>)	2	7%
Senegal Eremomela (<i>Eremomela pusilla</i>)	15	56%
Northern Crombec (<i>Sylvietta brachyura</i>)	5	19%

Database made by Turgor Henrik Dahl AB, SWEDEN

Species	Obs. days	Obs. %
Willow Warbler (Phylloscopus trochilus)	20	74%
Wood Warbler (Phylloscopus sibilatrix)	1	4%
Garden Warbler (Sylvia borin)	14	52%
Blackcap (Sylvia atricapilla)	2	7%
Common Whitethroat (Sylvia communis)	25	93%
Black-backed Cisticola (Cisticola eximius)	15	56%
Zitting Cisticola (Cisticola juncidis)	2	7%
Dorst's Cisticola (Cisticola dorsti)	5	19%
Rock-loving Cisticola (Cisticola aberrans)	21	78%
Singing Cisticola (Cisticola cantans)	21	78%
Winding Cisticola (Cisticola galactotes)	1	4%
Tawny-flanked Prinia (Prinia subflava)	24	89%
Red-winged Warbler (Heliolais erythroptera)	10	37%
Grey-backed Camaroptera (Camaroptera brachyura)	24	89%
Oriole-Warbler (Hypergerus atriceps)	10	37%
Pale Flycatcher (Bradornis pallidus)	2	7%
Western Black Flycatcher (Melaenornis edolioides)	22	81%
Pied Flycatcher (Ficedula hypoleuca)	16	59%
Spotted Flycatcher (Muscicapa striata)	4	15%
Senegal Batis (Batis senegalensis)	10	37%
Scarlet-spectacled Wattle-eye (Platysteira cyanea)	21	78%
Blue Flycatcher (Elminia longicauda)	8	30%
Paradise Flycatcher (Terpsiphone viridis)	11	41%
Brown Babbler (Turdoides plebejus)	3	11%
West African Penduline Tit (Anthoscopus parvulus)	1	4%
Pygmy Sunbird (Anthreptes platurus)	9	33%
Green-headed Sunbird (Nectarinia verticalis)	11	41%
Scarlet-chested Sunbird (Nectarinia senegalensis)	25	93%
Variable Sunbird (Nectarinia venusta)	24	89%
Coppery Sunbird (Nectarinia cuprea)	3	11%
Beautiful Sunbird (Nectarinia pulchella)	2	7%
Yellow White-eye (Zosterops senegalensis)	25	93%
African Golden Oriole (Oriolus auratus)	1	4%
Emin's Shrike (Lanius gubernator)	1	4%
Woodchat Shrike (Lanius senator)	1	4%

Database made by Turgor Henrik Dahl AB, SWEDEN

Species	Obs. days	Obs. %
Yellow-billed Shrike (<i>Corvinella corvina</i>)	5	19%
Northern Puffback (<i>Dryoscopus gambensis</i>)	1	4%
Black-crowned Tchagra (<i>Tchagra senegala</i>)	25	93%
Tropical Boubou (<i>Laniarius aethiopicus</i>)	22	81%
Yellow-crowned Gonolek (<i>Laniarius barbarus</i>)	25	93%
Sulphur-breasted Bush Shrike (<i>Malaconotus</i>	25	93%
Grey-headed Bush Shrike (<i>Malaconotus blanchoti</i>)	10	37%
Pied Crow (<i>Corvus albus</i>)	4	15%
Neumann's Starling (<i>Onychognathus morio</i>)	17	63%
Purple Glossy Starling (<i>Lamprotornis purpureus</i>)	24	89%
Lesser Blue-eared Starling (<i>Lamprotornis chloropterus</i>	3	11%
Splendid Glossy Starling (<i>Lamprotornis splendidus</i>)	7	26%
Northern Long-tailed Starling (<i>Lamprotornis caudatus</i>)	1	4%
Violet-backed Starling (<i>Cinnyricinclus leucogaster</i>)	25	93%
Yellow-billed Oxpecker (<i>Buphagus africanus</i>)	2	7%
Grey-headed Sparrow (<i>Passer griseus</i>)	22	81%
Bush Petronia (<i>Petronia dentata</i>)	5	19%
Speckle-fronted Weaver (<i>Sporopipes frontalis</i>)	25	93%
Chestnut-crowned Sparrow-weaver (<i>Plocepasser</i>	5	19%
Little Weaver (<i>Ploceus luteolus</i>)	11	41%
Black-necked Weaver (<i>Ploceus nigricollis</i>)	14	52%
Vitelline Masked Weaver (<i>Ploceus velatus</i>)	6	22%
Heuglin's Masked Weaver (<i>Ploceus heuglini</i>)	3	11%
Village Weaver (<i>Ploceus cucullatus</i>)	25	93%
Red-billed Quelea (<i>Quelea quelea</i>)	3	11%
Black-winged Red Bishop (<i>Euplectes hordeaceus</i>)	19	70%
Northern Red Bishop (<i>Euplectes franciscanus</i>)	20	74%
Yellow-mantled Whydah (<i>Euplectes macrourus</i>)	2	7%
White-cheeked Olive-back (<i>Nesocharis capistrata</i>)	1	4%
Bar-breasted Firefinch (<i>Lagonosticta rufopicta</i>)	2	7%
Red-billed Firefinch (<i>Lagonosticta senegala</i>)	25	93%
Black-bellied Firefinch (<i>Lagonosticta rara</i>)	4	15%
Rock Firefinch (<i>Lagonosticta sanguinodorsalis</i>)	25	93%
Lavender Waxbill (<i>Estrilda caerulescens</i>)	23	85%
Orange-cheeked Waxbill (<i>Estrilda melpoda</i>)	4	15%

Database made by Turgor Henrik Dahl AB, SWEDEN

Species	Obs. days	Obs. %
Black-rumped Waxbill (<i>Estrilda troglodytes</i>)	17	63%
Red-cheeked Cordon-bleu (<i>Uraeginthus bengalus</i>)	25	93%
Quail Finch (<i>Ortygospiza atricollis</i>)	17	63%
Warbling Silverbill (<i>Lonchura cantans</i>)	5	19%
Bronze Mannikin (<i>Lonchura cucullata</i>)	25	93%
Cut-throat Finch (<i>Amadina fasciata</i>)	3	11%
Village Indigobird (<i>Vidua chalybeata</i>)	3	11%
Pin-tailed Widow (<i>Vidua macroura</i>)	1	4%
White-rumped Seed-eater (<i>Serinus leucopygius</i>)	4	15%
Yellow-fronted Canary (<i>Serinus mozambicus</i>)	1	4%
Cinnamon-breasted Rock Bunting (<i>Emberiza tahapisi</i>)	25	93%
Number of	219	